

MEDICATION INFORMATION

*Use weight to determine dose especially if child was premature or is small for his/her age.

MEDICATIONS

| Age | 1-3 months | 4-6 months | 7-12 months | 13-24 months | 2-3 years |
|---|---------------|---------------|--------------------|--------------------|-------------------------------|
| Weight | 7-12 lbs | 13-18 lbs | 19-24 lbs | 25-30 lbs | 31-38 lbs |
| | 4 ml | 1/2 tsp | 3/4 tsp | 1 tsp | 1 1/4 tsp |
| | 1/4 tsp | 1/2 tsp | 3/4 tsp | 1 tsp | 1 1/4 tsp |
| Acetaminophen (Tylenol, Tempra, etc.) Every 4 hours as needed For fever or pain. Drops In ml, elixir in tsp. and Chewable in mgs. Instead | 4 ml | 8 ml | 1.2 ml | 1.6 ml | 2.0 ml |
| | 1/4 tsp | 1/2 tsp | 3/4 tsp | 1 tsp | 1 1/4 tsp |
| | | | | | <u>200 mgs</u> |
| Ibuprofen (Advil, Motrin) Every 6 hours, may be given | | | 3/4 tsp | 1 tsp | <u>2 1/2 mgs</u> 1 1/4 tsp |
| Pedicare infant drops Triaminic infant or Pediatric Sudafed Drops Every 6 hours as needed For nasal congestion | (check w/dr.) | 8 ml | 1.2 ml | 1.6 ml | 2.0 ml |
| Benadryl Elixir Every 4-6 hours for Allergic symptoms | (check w/dr.) | (check w/dr.) | 1/2 tsp | 1 tsp | 1 1/2 tsp |
| Dimetapp Elixir Triaminic Syrup (orange) Every 6 hours as needed For congestion and Drainage. May help Some types of coughs | (check w/dr.) | (check w/dr.) | 1/4 tsp 1/2 tsp | 1/2 tsp 3/4 tsp | 3/4 tsp 1 tsp |
| Robitussin or Robitussin PE may Be used for cough every 6 hours as needed | (check w/dr.) | (check w/dr.) | 1/4 tsp | 1/2 tsp | 3/4 tsp |

Physician's Signature

Date